SUPPORTING BREASTFEEDING


Fighting Childhood Obesity

Kiran Saluja MPH, RD



Celebrating WIC:
40 Years of Strengthening Families Through Advocacy





WIC's come a long way....40 years later



1989: First set aside for BF: \$8 mill

1992: Special Food package for bf women

1994: \$21 for every Pg and BF woman for BF

1997: Loving Support Launched

1998: \$24.80 for every Pg and BF woman

2004:\$20 million for Peer Counseling

2009: Food Packages changes to incentivize

exclusive breastfeeding

2014: \$60 million for Peer Counseling

THE REW WIC WORLD...



The WIC Program has stepped up to become the face of breastfeeding support in the community

- Active in coalitions/task forces
- Funds for peer counseling
- Terrific staff training curriculum
- NWA BF summit/strategic plan

ONLY PROGRAM WITH \$\$\$\$\$ to promote and support breastfeeding!!

....and we collect tons of data!

- WIC data: Proof that WIC works
- WIC can successfully promote breastfeeding
- Breastfeeding significantly reduces rates of childhood obesity

Breastfeeding and Obesity

PHFE WIC serves over 300,000 participants/month


~5000 births/month




- 60 sites in LA, Orange and San Bernardino Counties
- 84% Hispanic, 6% Asian, 5% African-American,
 4% White




















Breastfeeding reduces childhood obesity

- What WIC data do we have to support this?
 - 80,000+ infants on WIC from birth to 4+ years old
 - They have the same Individual ID for four + years



Initiation of Full breastfeeding = 23.6% reduction in obesity at age 4

Obesity Rate at Age 4



^{*}Package issuance at infant enrollment into WIC

Longer breastfeeding duration = lower obesity rates at age 4

- For every additional month of ANY breastfeeding:
 - 2% reduction in obesity at age 4.

- For every additional month of FULLY breastfeeding:
 - 4% reduction in obesity at age 4.

Breastfeeding, SSBs and Obesity: How are they related?

Study Goal:



To examine the separate and combined effects of breastfeeding in the first year of life and SSB intake at ages 2-4 on the prevalence of obesity in early childhood.



obesity prevalence only in children who were breastfed less than 12 months.

Davis, Whaley & Goran (2012). Effects of breastfeeding and low SSB intake on Hispanic toddlers. *Am J Clin Nut*, 95, 3-8. Funded by First 5 LA.

One out of two babies in the US is a WIC baby

















The Burden of Suboptimal
Breastfeeding in the United States
A Pediatric Cost Analysis
Melissa Bartick, MD, MSc, Arnold
Reinhold, MBA

PEDIATRICS (doi:10.1542/peds.2009-1616)


If 90% of US families could comply with medical recommendations to breastfeed exclusively for 6 months, the United States would save

\$13 billion per year

and prevent an excess


911 deaths,

nearly all of which would be in infants (\$10.5 billion and 741 deaths at 80% compliance).





By SABRINA TAVERNISE FEB. 25, 2014



heart disease and stroke.

The drop emerged from a major







WIC Breastfeeding Efforts Scaling New Heights

